

The 600 Ton ICARUS detector at the LNGS

David B. Cline

UCLA Astroparticle Physics Group

Topics

1. R & D on ICARUS technology detector method
2. Installation of the T600 at the LNGS
3. Physics program: (a) high energy neutrons from CERN, (b) Search for exotic proton decay
4. Possible future plans to search for sterile neutrinos
5. As a prototype for multi-kiloton detector around the world

Summary

The ICARUS Collaboration

M. Antonello, P. Aprili, N. Canci, C. Rubbia, E. Scantamburlo, E. Segreto, C. Vignoli
Laboratori Nazionali del Gran Sasso dell'INFN, Assergi (AQ), Italy

B. Baibussinov, M. BaldoCeolin, S. Centro, D. Dequal, C. Farnese, A. Fava, D. Gibin, A. Guglielmi, G. Meng, F. Pietropaolo, F. Varanini, S. Ventura
Dipartimento di Fisica e INFN, Università di Padova, Via Marzolo 8, I-35131, Padova, Italy

P. Benetti, E. Calligarich, R. Dolfini, A. Gigli Berzolari, A. Menegolli, C. Montanari, A. Rappoldi, G. L. Raselli, M. Rossella
Dipartimento di Fisica Nucleare e Teorica e INFN, Università di Pavia, Via Bassi 6, I-27100, Pavia Italy

F. Carbonara, A. G. Cocco, G. Fiorillo
Dipartimento di Scienze Fisiche, INFN e Università Federico II, Napoli, Italy

A. Cesana, P. Sala, A. Scaramelli, M. Terrani
INFN, Sezione di Milano e Politecnico, Via Celoria 2, I-20123

K. Cieslik, A. Dabrowska, M. Haranczyk, D. Stefan, M. Szarska, T. Wachala, A. Zalewska
The Henryk Niewodniczanski, Institute of Nuclear Physics, Polish Academy of Science, Krakow, Poland

D. B. Cline, S. Otwinowski, H.-G. Wang, X. Yang
Department of Physics and Astronomy, University of California, Los Angeles, USA

A. Dermenev, S. Gninenko, M. Kirsanov
INR RAS, prospekt 60-letiya Oktyabrya 7a, Moscow 117312, Russia

A. Ferrari
CERN, Ch1211 Geneve 23, Switzerland

T. Golan, J. Sobczyk, J. Zmuda
Institute of Theoretical Physics, Wroclaw University, Wroclaw, Poland

J. Holeczek, J. Kisiel, I. Kochanek, S. Mania
Institute of Physics, University of Silesia, 12 Bankowa st., 40-007 Katowice, Poland

J. Lagoda, T. J. Palczewski, P. Przewlocki, J. Stepaniak, R. Sulej
A. Soltan Institute for Nuclear Studies, 05-400 Swierk/Otwock, Warszawa, Poland

G. Mannocchi, L. Periale, P. Picchi,
Laboratori Nazionali di Frascati (INFN), Via Fermi 40, I-00044, Italy


P. Plonski, K. Zaremba
Institute for Radioelectronics, Warsaw Univ. of Technology Pl. Politechniki 1, 00-661 Warsaw, Poland

F. Sergiampietri
Dipartimento di Fisica, Università di Pisa, Largo Bruno Pontecorvo 3, I-56127, Pisa, Italy

A powerful detection technique

The **Liquid Argon Time Projection Chamber** [C. Rubbia: CERN-EP/77-08 (1977)] first proposed to INFN in 1985 [ICARUS: INFN/AE-85/7] capable of providing a 3D imaging of any ionizing event ('electronic bubble chamber') with in addition:


- continuously sensitive, self triggering
- high granularity (~ 1 mm)
- excellent calorimetric properties



Electrons from ionizing track are drifted in LAr by E_{drift} . They traverse transparent wire arrays oriented in different directions where induction signals are recorded. Finally electron charge is collected by collection plane.

- **Key feature: LAr purity from electro-negative molecules (O_2 , H_2O , CO_2).**
- **Target: 0.1 ppb O_2 equivalent = 3 ms lifetime (4.5 m drift @ $E_{\text{drift}} = 500$ V/cm).**


The path to larger LAr detectors



LAr-TPC performance

□ Tracking device:

- precise event topology ($s_{x,y} \sim 1\text{mm}$, $s_z \sim 0.4\text{mm}$)
- μ momentum measurement via multiple scattering:
 $\Delta p/p \sim 10\text{-}15\%$ depending on track length and p





□ Measurement of local energy deposition

dE/dx :

- $e/\geq$ separation (2% X_0 sampling);
- particle ID by means of dE/dx vs range

□ Total energy reconstruction by charge integration:

- full sampling, homogeneous calorimeter with excellent accuracy for contained events



RESOLUTIONS

- Low energy electrons: $\Delta(E)/E = 11\% / \sqrt{E(\text{MeV})} + 2\%$
- Electromagnetic showers: $\Delta(E)/E = 3\% / \sqrt{E(\text{GeV})}$
- Hadron shower (pure LAr): $\Delta(E)/E \approx 30\% / \sqrt{E(\text{GeV})}$



Summary of LAr TPC performances

- Tracking device
 - Precise event topology
 - Momentum via multiple scattering
 - Measurement of local energy deposition
 - e / γ separation ($2\%X_0$ sampling)
 - Particle ID by means of dE/dx vs range
 - Total energy reconstruction of the events from charge integration
 - Full sampling, homogeneous calorimeter with excellent accuracy for contained events
- RESOLUTIONS**

Low energy electrons: $\sigma(E)/E = 11\% / \sqrt{E(\text{MeV})} + 2\%$

Electromagn. showers: $\sigma(E)/E = 3\% / \sqrt{E(\text{GeV})}$

Hadron shower (pure LAr): $\sigma(E)/E \approx 30\% / \sqrt{E(\text{GeV})}$




dE/dx
distribution
along a single
muon track

Electrons from muon decay

- Excellent resolution obtained from the measured decay electron spectrum (Michel parameter) from muon decays



electrons from μ decays



The key feature of LAr imaging: very long e-mobility


- The main technological challenge of the development of the cryogenic liquid Ar chamber is the capability of ensuring a sufficiently long free electron lifetime.
- Indeed the free electron path in a liquid is ≈ 600 times shorter than in a gas. For instance **10 ms** lifetime corresponds to a **30 ppt (t=trillion !)** of Oxygen equivalent.
- At 500v/cm, a 5m drift length corresponds to a drift time of 3.1 ms (1.6 m/ms).
- The intrinsic bubble size (rms diffusion) is given by
- The values for 5m drift are $\langle\sigma_D\rangle \approx 1.1$ mm and $\sigma_{\max} \approx 1.6$ mm, tiny with respect to the wire pitch (≥ 2 mm).

$$\sigma_D [mm] = 0.9 \sqrt{T_D [ms]}$$





electron - π^0 identification



Reconstruction of π^0 -showers



Electron shower events are extremely well identified experimentally, because of the ionization behaviour in the first cells after the vertex.



The ICARUS T600 detector



■ Two identical modules

- $3.6 \times 3.9 \times 19.6 \approx 275 \text{ m}^3$ each
- Liquid Ar active mass: $\approx 476 \text{ t}$
- Drift length = 1.5 m
- HV = -75 kV E = 0.5 kV/cm

■ 4 wire chambers:

- 2 chambers per module
- 3 readout wire planes per chamber, wires at 0, $\pm 60^\circ$
- ≈ 54000 wires, 3 mm pitch, 3 mm plane spacing

■ PMT for scintillation light:

- (20+54) PMTs, 8" $\varnothing$


Venezia, 15-20/2011 $v_{drift} = 1.55 \text{ mm}/\mu\text{s}$

XIV Int. Workshop on Neutrino Telescopes



■ VUV sensitive (128nm) with wave shifter (TPB)

Slide: 8

ICARUS T600 in LNGS Hall B



LAr purity time evolution



Simple model: uniform distribution of the impurities, including internal degassing, decreasing in time, constant external leak and liquid purification by recirculation.

$$dN/dt = -N/\tau_R + k + k_I \exp(-t/\tau_I)$$

$$\tau_{ele} [\text{ms}] = 0.3 / N[\text{ppb O}_2 \text{ equivalent}]$$


τ_R : 2 m³/h corresponding to $\approx$ 6 day cycle time

τ_R : recirculation time for a full detector volume
 k_I and τ_I : related to the total degassing internal rate
 k : related to the external leaks

ICARUS T600 physics potential

- ❑ ICARUS T600: **major milestone** towards realization of large scale LAr detector. Interesting physics in itself: unique imaging capability, spatial/calorimetric resolutions and e/π^0 separation → **events “seen in a new Bubble chamber like” way.**
- ❑ CNGS ν events collection (beam intensity $4.5 \cdot 10^{19}$ pot/year, $E_\nu \sim 17.4$ GeV):
 - 1200 ν_μ CC event/year;
 - ~ 8 ν_e CC event/year;
 - observation of ν_τ events in the electron channel, using kinematical criteria;
 - search for sterile ν in LSND parameter space (deep inelastic ν_e CC events excess).
- ❑ “Self triggered” events collection:
 - ~ 80 events/y of unbiased atmospheric ν CC;
 - zero background proton decay with 3×10^{32} nucleons for “exotic” channels.

ICARUS and Status of Liquid Argon Technology



CNGS neutrino interactions in ICARUS T600

Drift time coordinate (1.4 m)


Collection view

Wire coordinate (8 m)

CNGS ν beam direction




ν_{μ} CC



ICARUS 9839/1274

CERN beam →



Candidate for a ν_e interaction

Preliminary results of first CNGS 2010 run

- Analyzed sample: 1494 CNGS triggers, i.e. $4.54 \cdot 10^{18}$ pot = 78 % out of whole sample. Classified by visual scanning into fiducial volume of 434 t.
- Number of collected interactions compared with number of interactions predicted ($(2.6 \text{ }_{-CC} + 0.86 \text{ }_{-NC}) 10^{-17}/\text{pot}$), in the whole energy range up to 100 GeV, corrected by fiducial volume and DAQ dead-time.


Event type	Collected	Expected
<u>CC</u>	94	98
<u>NC</u>	32	31
<u>XC</u> *	6	-
Total	132	129

* Events at edges, with μ track too short to be visually recognized: further analysis needed.

On overall statistics **in agreement with expectations.**

LAr-TPC: powerful technique. Run 9927 Event 572

12.5 m





Primary vertex (A)


very long μ (1),
e.m. cascade (2),
pion (3).

Secondary vertex (B)

The longest track (5) is a μ coming from stopping k (6). - μ decay is observed.




$$M_W^* = 125 \pm 15 \text{ MeV}/c^2$$



Track	$E_{\text{depl}} [\text{MeV}]$	$\cos x$	$\cos y$	$\cos z$
1 (μ)	2701.97	0.069	-0.040	-0.997
2 (π^0)	520.82	0.054	-0.420	-0.906
3 (π)	514.04	-0.001	0.137	-0.991
Sec. vtx.	797.			
4	76.99	0.009	-0.649	0.761
5 (μ)	313.9			
6 (K)	86.98	0.000	-0.239	-0.971
7	35.87	0.414	0.793	-0.446
8	283.28	-0.613	0.150	-0.776

Sterile neutrino search with ICARUS T600




$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ appearance search in T600 in LNSD parameter space

- Sensitivity region, in terms of standard deviations, for 3000 raw CNGS muon neutrino events.
- The potential signal is above the background generated by the intrinsic beam contamination, in the deep inelastic interval 10-30 GeV.
- Largely complementary to the Fermi-lab program in terms of energy and baseline.

ν_e CC interaction at ~ 1.5 GeV

- At these energies, electron identification and energy reconstruction of ν_e interactions is ensured with $\sim 5 X_0$ ($X_0=14\text{cm}$) longitudinal cut and $\sim 2 X_0$ side cut of the sensitive volume are performed, corresponding to a fiducial volume of $\sim 80\%$ of the active one.




π^0 from NC are rejected by photon vertex identification, invariant mass reconstruction and dE/dx measurement: the expected π^0 mis-interpretation probability is 0.1 %, with ν_e detection efficiency of 90 % within the fiducial volume.

With these fiducial cuts, the expected ν_e energy resolution is around 14 %

Two LAr-TPC detectors at the CERN-PS neutrino beam



CERN-SPSC/99-26
SPSC/P311
August 30, 1999

SEARCH FOR $\nu_\mu \rightarrow \nu_e$ OSCILLATION
AT THE CERN PS



Two positions are foreseen for the detection of the neutrinos
The far (ICARUS-T600) location at 850 m from the target: $L/E \sim 1 \text{ km/GeV}$;
The additional detector and new location at a distance of 127 m from the target: $L/E \sim 0.15 \text{ km/GeV}$

Comparing LSND sensitivities (*arXiv:0909.0355*)



Expected sensitivity for the proposed experiment exposed at the CERN-PS neutrino beam (left) for $2.5 \cdot 10^{20}$ pot (30 kW basic option) and twice as much for anti-neutrino (right). The LSND allowed region is fully explored both for neutrinos and anti-neutrinos. The T600 expectations from one year at LNGS are also shown.

Nucleon decay expectations

- LAr-TPC provides a much more powerful bkg rejection w.r.t. other techniques. It can perform a large variety of exclusive decay modes measurements in bkg free mode.
- With $1.2 \cdot 10^{34}$ nucleons, MODULAR is well suited for channels not accessible to — detectors due to the complicated event topology, or if the emitted particles are below the — threshold (e.g. K^+)

Channel	90% CL-5y	(pdg 90% CL)
$p \rightarrow \nu \pi^+$	$4.4 \cdot 10^{33}$	$(2.5 \cdot 10^{31})$
$p \rightarrow \mu^+ \pi^+ K^+$	$1.1 \cdot 10^{34}$	$(2.5 \cdot 10^{32})$
$n \rightarrow e^- K^+$	$1.3 \cdot 10^{34}$	$(3.2 \cdot 10^{31})$
$n \rightarrow \mu^+ \pi^-$	$6.0 \cdot 10^{33}$	$(1.0 \cdot 10^{32})$
$n \rightarrow \nu \pi^0$	$4.4 \cdot 10^{33}$	$(1.1 \cdot 10^{32})$

